

DATE/TIME	CHK'D BY	PHYSICIAN'S ORDER SHEET	
		ICU Propofol (DIPRIVAN®) PROTOCOL	Pharmacy
		Order Set No. 4090	
		Allergies:	Reactions:
		Allergies:	Reactions:
		Allergies:	Reactions:
		Attending M.D.:/DO/DDS	
		<input type="checkbox"/> Inpatient Admission <input type="checkbox"/> Outpatient <input type="checkbox"/> Observation <input type="checkbox"/> Extended Stay	
		Diagnosis:	
		Initiation and Maintenance of ICU sedation in mechanically ventilated patients with a secured airway:	
		Propofol (Diprivan®) can cause sudden and profound respiratory depression with minimal doses. Use with caution in patients with sepsis or severe acute pancreatitis	
		Start Propofol (Diprivan®) drip at 5 mcg/kg/min for at least 5 minutes Increase rate in 5-10 mcg/kg/min increments over 5-10 min until desired sedation MAX dose= 67 mcg/kg/min Change sedation agent at 48 hours	
		<input type="checkbox"/> Fentanyl drip 2000 mcg/200ml Start at 75mcg/hour IV titrate to MAAS of 2-3 <input checked="" type="checkbox"/> Triglyceride level at initiation and Q 3 days	
		Assess CNS function daily for sedation level Tubing and unused portions should be discarded after 12 hours Monitor patients for respiratory and circulatory depression Propofol (Diprivan®) contains 10 gm fat/100 ml; consider alternate agent if TG levels >300 mg/dl.	
		Phok42013apb	
		Page 1 of 1	
		Reviewed: April 2013	Physician's Signature

PROHIBITED ABBREVIATIONS			
PROHIBITED:	INSTEAD WRITE	PROHIBITED:	INSTEAD WRITE
MS, MgSO ₄	magnesium sulfate	Q.D.	daily
MS, MSO ₄	morphine sulfate	QOD	every other day
U	units	.5 mg	0.5 mg – always use zero before decimal
IU	international units	5.0 mg	5 mg – never use trailing zeroes

Patient Identification